

Types of Biological Samples Collected at Follow-Up 1

Type	Volume (mL)	Number	Total volume (mL)	Number of aliquots
1. Citrate (<i>3.2% buffered sodium citrate solution, 0.109 M</i>)				
Plasma	2.7	1	2.7	2
Plasma, platelet poor	2.7	1	2.7	2
2. Serum (<i>spray coated silicone and micronized silica particles</i>)				
Serum	10	1	10	8
3. Heparin (<i>lithium heparin, 90 USP</i>)				
Plasma	10	1	10	6
4. EDTA (<i>10.8 mg spray coated K₂ EDTA</i>)				
Plasma	3	1	3	0 ¹
Plasma	6	3	18	6
Buffy coat			(2)	4
Whole blood			(1.0)	2
PBMC ²	6	1	6	3 ³ 3 ⁴
5. ACD (<i>13.2 g/L trisodium citrate, 4.8 g/L citric acid, and 14.7 g/L dextrose; 0.4 mL</i>)				
Whole blood	6	1	6.0	2
6. Urine				
Urine	60 ⁵	1	2	4

¹ Sample is used for performing hematology tests at the DCS.

² Obtained using the Greiner LeucoSep tube to produce peripheral blood mononuclear cells (PBMC).

³ With PBS.

⁴ With cell culture media (RPMI 1640 Medium with 12.5% human serum albumin and 12.5% High Grade DMSO).

⁵ Random collection, variable volume.